arXiv:0910.0692v1 [math.DG] 5 Oct 2009

BIHARMONIC MAPS INTO COMPACT LIE GROUPS AND THE INTEGRABLE SYSTEMS

HAJIME URAKAWA

Abstract. In this paper, the reduction of biharmonic map equation in terms of the Maurer-Cartan form for all smooth map of an arbitrary compact Riemannian manifold into a compact Lie group (G,h) with bi-invariant Riemannian metric h is obtained. By this formula, all biharmonic curves into compaqct Lie groups are determined, and all the biharmonic maps of an open domain of $\mathbb{R}^2$ with the conformal metric of the standard Riemannian metric into (G,h) are determined.

1. Introduction and statement of results.

The theory of harmonic maps into Lie groups, symmetric spaces or homogeneous spaces has been extensively studied related to the integrable systems by many mathematicians (for examples, [6], [7], [3], [2]). In this paper, we want to build a bridge between the theory of biharmonic maps into Lie groups and the integrable systems extending the above harmonic map theory.

Acknowledgement: The author expresses his gratitude to Prof. J. Inoguchi who gave many useful suggestions and Prof. A. Kasue for his financial support during the preparation of this paper .

2. Preliminaries.

In this section, we prepare general materials and facts harmonic maps, biharmonic maps into compact Lie groups (cf. [5]).

Let (M,g) be an m dimensional compact Riemannian manifold, and G be an n dimensional compact Lie group with Lie algebra $\mathfrak{g}$, and h,

---

2000 *Mathematics Subject Classification.* 58E20.

Key words and phrases. harmonic map, biharmonic map, compact Lie group, integrable system, Maurer-Cartan form.

Supported by the Grant-in-Aid for the Scientific Research, (A), No. 19204004; (C), No. 21540207, Japan Society for the Promotion of Science.

the bi-invariant Riemannian metric on G corresponding to the $\mathrm{Ad}(G)$-invariant inner product $\langle\ ,\ \rangle$ on $\mathfrak{g}$.

The *energy functional* on the space $C^\infty(M,G)$ of all C^∞ maps of M into G is defined by

$$E(\psi)=\frac{1}{2}\int_M \|d\psi\|^2\,v_g,$$

and for a C^∞ one parameter deformation $\psi_t\in C^\infty(M,G)$ $(-\epsilon<t<\epsilon)$ of ψ with $\psi_0=\psi$, the *first variation formula* is given by

$$\left.\frac{d}{dt}\right|_{t=0} E(\psi_t)=-\int_M\langle\tau(\psi),V\rangle\,v_g,$$

where V is a variation vector field along ψ defined by $V=\left.\frac{d}{dt}\right|_{t=0}\psi_t$ which belongs to the space $\Gamma(\psi^{-1}TG)$ of sections of the induced bundle of the tangent bundle TG by ψ. The *tension field* $\tau(\psi)$ is defined by

$$\tau(\psi)=\sum_{i=1}^m B(\psi)(e_i,e_i),\tag{2.1}$$

where

$$B(\psi)(X,Y)=\nabla^h_{d\psi(X)}d\psi(Y)-d\psi(\nabla_X Y)$$

for $X,Y\in\mathfrak{X}(M)$. Here, ∇, and ∇^h, are the Levi-Civita connections of (M,g) and (G,h), respectively. For a harmonic map $\psi:(M,g)\to(G,h)$, the *second variation formula* of the energy functional $E(\psi)$ is

$$\left.\frac{d^2}{dt^2}\right|_{t=0} E(\psi_t)=\int_M\langle J(V),V\rangle\,g_g$$

where

$$\begin{aligned}J(V)&=\overline{\Delta}V-\mathcal{R}(V),\\ \overline{\Delta}V&=\overline{\nabla}^*\,\overline{\nabla}V=-\sum_{i=1}^m\{\overline{\nabla}_{e_i}(\overline{\nabla}_{e_i}V)-\overline{\nabla}_{\nabla_{e_i}e_i}V\},\\ \mathcal{R}(V)&=\sum_{i=1}^m R^h(V,d\psi(e_i))d\psi(e_i).\end{aligned}$$

Here, $\overline{\nabla}$ is the induced connection on the induced bundle $\psi^{-1}TG$, and is R^h is the curvature tensor of (G,h) given by $R^h(U,V)W=[\nabla^h_U,\nabla^h_V]W-\nabla^h_{[U,V]}W$ $(U,V,W\in\mathfrak{X}(G))$.

The *bienergy functional* is defined by

$$E_2(\psi)=\frac{1}{2}\int_M\|(d+\delta)^2\psi\|^2\,v_g=\frac{1}{2}\int_M|\tau(\psi)|^2\,v_g,\tag{2.2}$$

and the *first variation formula* of the bienergy is given (cf. [4]) by

$$\left.\frac{d}{dt}\right|_{t=0} E_2(\psi_t) = -\int_M \langle \tau_2(\psi), V\rangle \, v_g \tag{2.3}$$

where the *bitension field* $\tau_2(\psi)$ is defined by

$$\tau_2(\psi) = J(\tau(\psi)) = \overline{\Delta}\tau(\psi) - \mathcal{R}(\tau(\psi)), \tag{2.4}$$

and a C^∞ map $\psi : (M, g) \to (G, h)$ is called to be *biharmonic* if

$$\tau_2(\psi) = 0. \tag{2.5}$$

3. Determination of the bitension field

Now, let θ be the Maurer-Cartan form on G, i.e., a $\mathfrak{g}$-valued left invariant 1-form on G which is defined by $\theta_y(Z_y) = Z$, $(y \in G,\, Z \in \mathfrak{g})$. For every C^∞ map ψ of (M, g) into (G, h), let us consider a $\mathfrak{g}$-valued 1-form α on M given by $\alpha = \psi^*\theta$. Then, it is well known (see for example, [2]) that

Lemma 3.1. *For every C^∞ map $\psi : (M, g) \to (G, h)$,*

$$\theta(\tau(\psi)) = -\delta\alpha \tag{3.1}$$

where $\alpha = \psi^*\theta$, and θ is the Maurer-Cartan form of G.

Thus, $\psi : (M, g) \to (G, h)$ is harmonic if and only if $\delta\alpha = 0$.

Furthemore, let $\{X_s\}_{s=1}^n$ be an orthonormal basis of $\mathfrak{g}$ with respect to the inner product $\langle\, ,\, \rangle$. Then, for every $V \in \Gamma(\psi^{-1}TG)$,

$$\begin{aligned} V(x) &= \sum_{i=1}^n h_{\psi(x)}(V(x), X_{s\,\psi(x)})\, X_{s\,\psi(x)} \in T_{\psi(x)}G, \\ \theta(V)(x) &= \sum_{i=1}^n h_{\psi(x)}(V(x), X_{s\,\psi(x)})\, X_s \in \mathfrak{g}, \end{aligned} \tag{3.2}$$

for all $x \in M$. Then, for every $X \in \mathfrak{X}(M)$,

$$\begin{aligned} \theta(\overline{\nabla}_X V) &= \sum_{s=1}^n h(\overline{\nabla}_X V, X_s)\, X_s \\ &= \sum_{s=1}^n \{X\, h(V, X_s)\, X_s - h(V, \overline{\nabla}_X X_s)\} X_s \\ &= X(\theta(V)) - \sum_{s=1}^n h(V, \overline{\nabla}_X X_s) X_s, \end{aligned} \tag{3.3}$$

where we regarded a vector field $Y \in \mathfrak{X}(G)$ to be an element in $\Gamma^{-1}(TG)$ by

$$Y(x) = Y(\psi(x)), \quad (x \in M),$$

and we useed the compatibility condition:

$$X\, h(V,W) = h(\overline{\nabla}_X V, W) + h(V, \overline{\nabla}_X W), \quad (V, W \in \Gamma(\psi^{-1}TG)),$$

for all $X \in \mathfrak{X}(M)$.

For every $X \in \mathfrak{X}(M)$, the differential of ψ, $\psi_* X \in \Gamma(\psi^{-1}TG)$ which is given by $\psi_* X_x \in T_{\psi(x)}G$ $(x \in M)$, can be written as

$$\psi_* X_x = \sum_{t=1}^{n} h(\psi_* X_x, X_{t\,\psi(x)}) X_{t\,\psi(x)}, \quad (x \in M),$$

which yields that

$$(\overline{\nabla}_X X_s)_x = \sum_{t=1}^{n} h(\psi_* X_x, X_{t\,\psi(x)})\, (\nabla^h_{X_t} X_s)_{\psi(x)} \tag{3.4}$$

for all $x \in M$.

Here, let us recall that the Levi-Civita connection ∇^h of (G,h) is given (cf. [5] Vol. II, p. 201, Theorem 3.3) by

$$\nabla^h_{X_t} X_s = \frac{1}{2}\,[X_t, X_s] = \frac{1}{2} \sum_{\ell=1}^{n} C_{ts}^{\ell}\, X_\ell, \tag{3.5}$$

where the structure constant C_{ts}^{ℓ} of $\mathfrak{g}$ is defined by

$$[X_t, X_s] = \sum_{\ell=1}^{n} C_{ts}^{\ell}\, X_\ell,$$

and satisfies that

$$\begin{aligned} C_{ts}^{\ell} &= \langle [X_t, X_s], X_\ell \rangle \\ &= -\langle X_s, [X_t, X_\ell] \rangle \\ &= -C_{t\ell}^{s}. \end{aligned} \tag{3.6}$$

Thus, we have by (3.4) and (3.5),

$$\begin{aligned}\sum_{s=1}^{n} h(V, \overline{\nabla}_X X_s) X_s &= \frac{1}{2}\sum_{s,t=1}^{n} h\left(V, \sum_{\ell=1}^{n} h(\psi_* X, X_t) C_{ts}^{\ell}\, X_\ell\right) X_s \\ &= -\frac{1}{2}\sum_{s,t,\ell=1}^{n} h(V, X_\ell)\, h(\psi_* X, X_t)\, C_{t\ell}^{s} X_s \\ &= -\frac{1}{2}\sum_{t,\ell=1}^{n} h(V, X_\ell)\, h(\psi_* X, X_t)\, [X_t, X_\ell] \\ &= -\frac{1}{2}\left[\sum_{t=1}^{n} h(\psi_* X, X_t)\, X_t, \sum_{\ell=1}^{n} h(V, X_\ell)\, X_\ell\right] \\ &= -\frac{1}{2}\left[\alpha(X), \theta(V)\right], \qquad (3.7)\end{aligned}$$

because we have

$$\begin{aligned}\alpha(X) &= (\psi^*\theta)(X) \\ &= \theta(\psi_* X) \\ &= \theta\left(\sum_{t=1}^{n} h(\psi_* X, X_t) X_t\right) \\ &= \sum_{t=1}^{n} h(\psi_* X, X_t) X_t, \qquad (3.8)\end{aligned}$$

and

$$\begin{aligned}\theta(V) &= \sum_{\ell=1}^{n} h(V, X_\ell)\, \theta(X_\ell) \\ &= \sum_{\ell=1}^{n} h(V, X_\ell)\, X_\ell. \qquad (3.9)\end{aligned}$$

Substituting (3.8) and (3.9) into the above, we have the last equality of (3.7).

Therefore, we obtain the following lemma together with (3.7) and (3.3).

Lemma 3.2. *For every C^∞ map $\psi :\, (M, g) \to (G, h)$,*

$$\theta(\overline{\nabla}_X V) = X(\theta(V)) + \frac{1}{2}\,[\alpha(X), \theta(V)], \qquad (3.10)$$

where $V \in \Gamma(\psi^{-1}TG)$ and $X \in \mathfrak{X}(M)$.

□

We show

Theorem 3.3. *For every* $\psi \in C^{\infty}(M,G)$*, we have*

$$\begin{aligned}\theta(\tau_2(\psi)) &= \theta(J(\tau(\psi)))\\ &= -\delta\, d\, \delta\alpha - \mathrm{Trace}_g([\alpha, d\,\delta\alpha]),\end{aligned} \tag{3.11}$$

where $\alpha = \psi^{*}\theta$*.*

Here, let us recall the definition:

Definition 3.4. *For two* $\mathfrak{g}$*-valued* 1*-forms* α *and* β *on* M*, we define* $\mathfrak{g}$*-valued symmetric* 2*-tensor* $[\alpha,\beta]$ *on* M *by*

$$[\alpha,\beta](X,Y) := \frac{1}{2}\left\{[\alpha(X),\beta(Y)] - [\beta(X),\alpha(Y)]\right\}, \quad (X,Y \in \mathfrak{X}(M)) \tag{3.12}$$

and its trace $\mathrm{Trace}_g([\alpha,\beta])$ *by*

$$\mathrm{Trace}_g([\alpha,\beta]) := \sum_{i=1}^{m} [\alpha,\beta](e_i,e_i). \tag{3.13}$$

We also use a $\mathfrak{g}$*-valued* 2*-form* $[\alpha\wedge\beta]$ *on* M *by*

$$[\alpha\wedge\beta](X,Y) := \frac{1}{2}\left\{[\alpha(X),\beta(Y)] - [\alpha(Y),\beta(X)]\right\}, \quad (X,Y \in \mathfrak{X}(M)). \tag{3.14}$$

Then, we have imediately by Theorem 3.3,

Corollary 3.5. *For every* $\psi \in C^{\infty}(M,G)$*, we have*
(1) $\psi : (M,g) \to (G,h)$ *is harmonic if and only if*

$$\delta\alpha = 0. \tag{3.15}$$

(2) $\psi : (M,g) \to (G,h)$ *is biharmonic if and only if*

$$\delta\, d\, \delta\alpha + \mathrm{Trace}_g([\alpha, d\,\delta\alpha]) = 0. \tag{3.16}$$

We give a proof of Theorem 3.3.

Proof. (The first step) We first show that, for all $V \in \Gamma(\psi^{-1}TG)$,

$$\begin{aligned}\theta(\overline{\Delta}V) &= \Delta_g\theta(V)\\ &\quad - \sum_{i=1}^{m}\left\{\frac{1}{2}[e_i(\alpha(e_i)),\theta(V)] + [\alpha(e_i), e_i(\theta(V))]\right.\\ &\qquad\qquad + \frac{1}{4}[\alpha(e_i),[\alpha(e_i),\theta(V)]]\\ &\qquad\qquad\qquad \left. -\frac{1}{2}[\alpha(\nabla_{e_i}e_i),\theta(V)]\right\},\end{aligned} \tag{3.17}$$

where $\{e_i\}_{i=1}^m$ is a locally defined orthonormal frame field on (M, g), and Δ_g is the (positive) Laplacian of (M, g) acting on $C^\infty(M)$.

Indeed, we have by using Lemma 3.2 twice,

$$\begin{aligned}
\theta(\overline{\Delta}V) &= -\sum_{i=1}^m \Big\{\theta(\overline{\nabla}_{e_i}(\overline{\nabla}_{e_i}V)) - \theta(\overline{\nabla}_{\nabla_{e_i}e_i}V)\Big\} \\
&= -\sum_{i=1}^m \Big\{e_i(\theta(\overline{\nabla}_{e_i}V)) + \frac{1}{2}[\alpha(e_i), \theta(\overline{\nabla}_{e_i}V)] \\
&\qquad\qquad -\nabla_{e_i}e_i(\theta(V)) - \frac{1}{2}[\alpha(\nabla_{e_i}e_i), \theta(V)]\Big\} \\
&= -\sum_{i=1}^m \Big\{e_i\Big(e_i(\theta(V)) + \frac{1}{2}[\alpha(e_i), \theta(V)]\Big) \\
&\qquad\qquad + \frac{1}{2}\Big[\alpha(e_i), e_i(\theta(V)) + \frac{1}{2}[\alpha(e_i), \theta(V)]\Big] \\
&\qquad\qquad -\nabla_{e_i}e_i\,(\theta(V)) - \frac{1}{2}[\alpha(\nabla_{e_i}e_i), \theta(V)]\Big\} \\
&= -\sum_{i=1}^m \{e_i(e_i(\theta(V))) - \nabla_{e_i}e_i\,(\theta(V))\} \\
&\quad -\sum_{i=1}^m \Big\{\frac{1}{2}e_i([\alpha(e_i), \theta(V)]) + \frac{1}{2}[\alpha(e_i), e_i(\theta(V))] \\
&\qquad\qquad\qquad + \frac{1}{4}[\alpha(e_i), [\alpha(e_i), \theta(V)]] \\
&\qquad\qquad\qquad\quad -\frac{1}{2}[\alpha(\nabla_{e_i}e_i), \theta(V)]\Big\}. \qquad (3.18)
\end{aligned}$$

Here, we have

$$e_i([\alpha(e_i), \theta(V)]) = [e_i(\alpha(e_i)), \theta(V)] + [\alpha(e_i), e_i(\theta(V))],$$

which we substitute into (3.18), and by definition of Δ_g, we have (3.17).

(The second step) On the other hand, we have to consider

$$-\sum_{i=1}^m R^h(V, \psi_* e_i))\psi_* e_i = -\sum_{i=1}^m R^h(L_{\psi(x)\,*}^{-1}V, L_{\psi(x)\,*}^{-1}\psi_* e_i)L_{\psi(x)\,*}^{-1}\psi_* e_i. \qquad (3.19)$$

Under the identification $T_eG \ni Z_e \leftrightarrow Z \in \mathfrak{g}$, we have

$$T_eG \ni L_{\psi(x)\,*}^{-1}\psi_* e_i \leftrightarrow \alpha(e_i) \in \mathfrak{g}, \qquad (3.20)$$

$$T_eG \ni L_{\psi(x)\,*}^{-1}V \leftrightarrow \theta(V) \in \mathfrak{g}, \qquad (3.21)$$

respectively.

Because, we have

$$L_{\psi(x)\,*}^{-1}\psi_*e_i=\sum_{i=1}^n h(\psi_*e_i,X_{s\,\psi(x)})\,X_{s\,e}$$

and

$$\begin{aligned}\alpha(e_i)&=\psi^*\theta(e_i)\\&=\theta(\psi_*e_i)\\&=\sum_{s=1}^n h(\psi_*e_i,X_{s\,\psi(x)})\,\theta(X_{s\,\psi(x)})\\&=\sum_{s=1}^n h(\psi_*e_i,X_{s\,\psi(x)})\,X_s,\end{aligned}$$

which implies that (3.20). By the same way, we have (3.21).

Under this identification, the curvature tensor (G,h) is given as (see Kobayashi-Nomizu ([5], pp. 203–204)),

$$R^h(X,Y)_e=-\frac{1}{4}\mathrm{ad}([X,Y])\qquad(X,Y\in\mathfrak{g}),$$

and then, we have

$$\begin{aligned}\theta\left(-\sum_{i=1}^m R^h(V,\psi_*e_i))\psi_*e_i\right)&=\frac{1}{4}\sum_{i=1}^m\left[[\theta(V),\alpha(e_i)]\,,\alpha(e_i)\right]\\&=\frac{1}{4}\sum_{i=1}^m\left[\alpha(e_i),[\alpha(e_i),\theta(V)]\right].\end{aligned}\tag{3.22}$$

(The third step) By (3.17) and (3.22), for $V \in \Gamma(\psi^{-1}TG)$, we have

$$\begin{aligned}
&\theta\left(\overline{\Delta}V - \sum_{i=1}^m R^h(V, \psi_* e_i)\psi_* e_i\right) \\
&= \Delta_g \theta(V) \\
&\quad - \sum_{i=1}^m \Big\{ \frac{1}{2}[e_i(\alpha(e_i)), \theta(V)] + [\alpha(e_i), e_i(\theta(V))] \\
&\qquad\qquad + \frac{1}{4}[\alpha(e_i), [\alpha(e_i), \theta(V)]] \\
&\qquad\qquad\quad - \frac{1}{2}[\alpha(\nabla_{e_i} e_i), \theta(V)] \Big\} \\
&\quad + \frac{1}{4}\sum_{i=1}^m [\alpha(e_i), [\alpha(e_i), \theta(V)]] \\
&= \Delta_g \theta(V) - \frac{1}{2}\sum_{i=1}^m e_i(\alpha(e_i)), \theta(V)] + \sum_{i=1}^m [\alpha(e_i), e_i(\theta(V))] \\
&\qquad\qquad + \frac{1}{2}\sum_{i=1}^m [\alpha(\nabla_{e_i} e_i), \theta(V)] \\
&= \Delta_g \theta(V) - \frac{1}{2}\left[\sum_{i=1}^m \left(e_i(\alpha(e_i)) - \alpha(\nabla_{e_i} e_i)\right), \theta(V)\right] \\
&\qquad + \sum_{i=1}^m [\alpha(e_i), e_i(\theta(V))] \\
&= \Delta_g \theta(V) + \frac{1}{2}[\delta\alpha, \theta(V)] + \sum_{i=1}^m [\alpha(e_i), e_i(\theta(V))]. \qquad (3.23)
\end{aligned}$$

(The forth step) For $V = \tau(\psi)$ in (3.23), since $\theta(\tau(\psi)) = -\delta\alpha$, we have

$$\begin{aligned}
\theta(J(\tau(\psi))) &= \Delta_g(\theta(\tau(\psi)) + \frac{1}{2}[\delta\alpha, \theta(\tau(\psi))] \\
&\qquad + \sum_{i=1}^m [\alpha(e_i), e_i(\theta(\tau(\psi))] \\
&= -\Delta_g \delta\alpha - \frac{1}{2}[\delta\alpha, \delta\alpha] - \sum_{i=1}^m [\alpha(e_i), e_i(\delta\alpha)] \\
&= -\Delta_g \delta\alpha - \sum_{i=1}^m [\alpha(e_i), e_i(\delta\alpha)] \\
&= \Delta_g \delta\alpha - \sum_{i=1}^m [\alpha(e_i), d\delta\alpha(e_i)]. \qquad (3.24)
\end{aligned}$$

Then, (3.24) implies the desired (3.11). □

4. Biharmonic maps from $\mathbb{R}$ into compact Lie groups

In this section, we consider the simplest case: the standard 1-dimensional Euclidean space, $(M, g) = (\mathbb{R}, g_0)$ and (G, h), an n-dimensional compact Lie group with the bi-invariant Riemannian metric h.

4.1 First, let $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in (G, h)$, a C^∞ curve in G. Then, $\alpha := \psi^*\theta$ is a $\mathfrak{g}$-valued 1-form on $\mathbb{R}$. So, α can be written at $t \in \mathbb{R}$ as

$$\alpha_t = F(t)\, dt \tag{4.1}$$

where $F : \mathbb{R} \ni t \mapsto F(t) \in \mathfrak{g}$ is a C^∞ function on $\mathbb{R}$ given by

$$F(t) = \alpha\left(\frac{\partial}{\partial t}\right) = \psi^*\theta\left(\frac{\partial}{\partial t}\right) = \theta\left(\psi_*\left(\frac{\partial}{\partial t}\right)\right). \tag{4.2}$$

Here, since

$$\psi'(t) := \psi_*\left(\frac{\partial}{\partial t}\right) = \sum_{i=1}^n h_{\psi(t)}\left(\psi_*\left(\frac{\partial}{\partial t}\right), X_{s\,\psi(t)}\right) X_{s\,\psi(t)}, \tag{4.3}$$

we have

$$F(t) = \sum_{s=1}^n h_{\psi(t)}\left(\psi_*\left(\frac{\partial}{\partial t}\right), X_{s\,\psi(t)}\right) X_s, \tag{4.4}$$

so that we have the following correspondence:

$$\begin{aligned} T_eG \ni & L_{\psi(t)\,*}^{-1}\psi'(t) = \sum_{s=1}^n h_{\psi(t)}(\psi'(t), X_{s\,\psi(t)})\, X_{s\,e} \\ & \leftrightarrow F(t) = \theta\left(\psi_*\left(\frac{\partial}{\partial t}\right)\right) \in \mathfrak{g}. \end{aligned} \tag{4.5}$$

4.2 We first show that

$$\delta\alpha = -F'(t). \tag{4.6}$$

Proof. Indeed, we have

$$\begin{aligned} \delta\alpha &= -\{\nabla^h_{e_i}(\alpha(e_i)) - \alpha(\nabla_{e_1}e_1)\} \\ &= -e_1(\alpha(e_1)) \\ &= -e_1(F(t)) \\ &= -F'(t). \end{aligned}$$

□

Therefore, we have $\psi:\,(\mathbb{R},g_0)\to(G,h)$ is *harmonic* if and only if

$$\begin{aligned}\delta\alpha=0\quad&\Longleftrightarrow\quad F'=0\\&\Longleftrightarrow\quad \alpha=X\otimes dt\qquad(\exists\ X\in\mathfrak{g})\\&\Longleftrightarrow\quad \psi:\,\mathbb{R}\to(G,h),\ \textit{geodesic},\end{aligned}\tag{4.7}$$

since

$$F(t)=\theta(\psi'(t))=L_{\psi(t)*}{}^{-1}\psi'(t),\tag{4.8}$$

we have

$$\psi'(t)=L_{\psi(t)*}X,\tag{4.9}$$

for some $X\in\mathfrak{g}$ which yields that

$$\psi(t)=x\,\exp(tX+Y)$$

for some $Y\in\mathfrak{g}$.

Therefore, *any geodesic through* $\psi(0)=x$ *is given by*

$$\psi(t)=x\,\exp(tX+Y),\quad (t\in\mathbb{R})\tag{4.10}$$

for some $X\in\mathfrak{g}$ and $Y\in\mathfrak{g}$.

On the other hand, we want to determine a *biharmonic curve* $\psi:\,(\mathbb{R},g_0)\to(G,h)$. By (4.6), we have

$$\delta d\delta\alpha=-\frac{\partial^2}{\partial t^2}\left(-F'(t)\right)=F^{(3)}(t),\tag{4.11}$$

and

$$\mathrm{Trace}_g[\alpha,d\delta\alpha]=\left[\alpha\left(\frac{\partial}{\partial t}\right),\,d\delta\alpha\left(\frac{\partial}{\partial t}\right)\right]=\left[F(t),F''(t)\right],\tag{4.12}$$

so by (4.9), (4.10), and (3.16) in Corollary 3.5, $\psi:\,(\mathbb{R},g_0)\to(G,h)$ is *biharmonic* if and only if

$$F^{(3)}-[F(t),F''(t)]=0.\tag{4.13}$$

By choosing a local coordinate on a neighborhood U around $e\in G$ as

$$U\ni\exp\left(\sum_{i=1}^n x_i\,X_i\right)\mapsto(x_1,\cdots,x_n)\in\mathbb{R}^n,$$

we may take as a local coordinate on $x\,U$ around each $x\in G$

$$x\,U\ni x\,\exp\left(\sum_{i=1}^n x_i\,X_i\right)\mapsto(x_1,\cdots,x_n)\in\mathbb{R}^n.$$

Then, every C^∞ curve $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in G$ with $\psi(0) = x$ can be written locally by

$$\psi(t) = x \exp\left(\sum_{i=1}^{n} x_i(t)\, X_i\right), \qquad (t \in \mathbb{R}).$$

Then, for (4.8),

$$F(t) = \sum_{i=1}^{n} h_{\psi(t)}(\psi'(t), X_{i\,\psi(t)} X_i) \tag{4.14}$$

since

$$L_{\psi(t)*}{}^{-1}\psi'(t) = \sum_{i=1}^{n} h_{\psi(t)}(\psi'(t), X_i\, \psi(t))\, X_{i\,e} \in T_eG. \tag{4.15}$$

and for (4.15),

$$L_{\psi(t)*}{}^{-1}\psi'(t) = \sum_{i=1}^{n} x_i{}'(t)\, X_i, \tag{4.16}$$

so we have

$$F(t) = \sum_{i=1}^{n} x_i{}'(t)\, X_i. \tag{4.17}$$

Thus, a C^∞ curve $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in G$ is *biharmonic* if and only if

$$x_k^{(4)}(t) - \sum_{i,j=1}^{n} x_i'(t)\, x_j^{(3)}(t)\, C_{ij}^k = 0 \quad (k = 1, \cdots, n), \tag{4.18}$$

where we put $[X_i, X_j] = \sum_{k=1}^n C_{ij}^k\, X_k$. Therefore, if we put $y_k(t) := x_k'(t)$ $(k = 1, \cdots, n)$, (4.13) is reduced to the following third order ODE's:

$$y_k^{(3)}(t) - \sum_{i,j=1}^{n} y_i(t)\, y_j''(t)\, C_{ij}^k = 0 \quad (k = 1, \cdots, n). \tag{4.19}$$

Thsu, we can summarize the above as follows:

Theorem 4.1. (1) *A C^∞ curve $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in (G, h)$ is harmonic if and only if $\psi(t) = x\, \exp(t\, X + Y)$ for some X, $Y \in \mathfrak{g}$.*

(2) *A C^∞ curve $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in (G, h)$ is biharmonic if and only if the ODE 's* (4.18) *or* (4.19) *hold.*

4.3 Now, we determine a *biharmonic* curve $\psi : \mathbb{R} \ni t \mapsto \psi(t) \in (G, h)$ when $G = SU(2)$ with the biinvariant Riemannian metric h

which corresponds to the following $\mathrm{Ad}(SU(2))$-invariant inner product $\langle\ ,\ \rangle$ on

$$\begin{aligned}&\mathfrak{g} = \mathfrak{su}(2) = \{X \in M(2,\mathbb{C});\ X + {}^{\mathrm{t}}\overline{X} = 0, \mathrm{Tr}(X) = 0\}.\\ &\langle X, Y\rangle = -2\mathrm{Tr}(XY) \quad (X, Y \in \mathfrak{su}(2)).\end{aligned}$$

If we choose

$$X_1 = \begin{pmatrix} \frac{\sqrt{-1}}{2} & 0 \\ 0 & -\frac{\sqrt{-1}}{2} \end{pmatrix},\ X_2 = \begin{pmatrix} 0 & \frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix},\ X_3 = \begin{pmatrix} 0 & \frac{\sqrt{-1}}{2} \\ \frac{\sqrt{-1}}{2} & 0 \end{pmatrix},$$

then $\{X_1, X_2, X_3\}$ is an orthonormal basis of $(\mathfrak{su}(2), \langle\ ,\ \rangle)$, and satisfies the Lie bracket relations:

$$[X_1, X_2] = X_3,\ [X_2, X_3] = X_1,\ [X_3, X_1] = X_2.$$

Thus, the ODE's (4.19) becomes

$$\begin{cases} y_1^{(3)} = y_2\, y_3'' - y_3\, y_2'', \\ y_2^{(3)} = y_3\, y_1'' - y_1\, y_3'', \\ y_3^{(3)} = y_1\, y_2'' - y_2\, y_1'', \end{cases} \tag{4.20}$$

which is equivalent to

$$\mathbf{y}^{(3)} = \mathbf{y} \times \mathbf{y}'', \tag{4.21}$$

where $\mathbf{y} := {}^{\mathrm{t}}(y_1, y_2, y_3) \in \mathbb{R}^3$, and $\mathbf{a} \times \mathbf{b}$ stands for the cross product in $\mathbb{R}^3$. The ODE's (4.20) can be solved as follows.

Let $\mathbf{x}(s) = {}^{\mathrm{t}}(x_1(s), x_2(s), x_3(s))$ be a C^∞ curve in $\mathbb{R}^3$ with arc length parameter s, and then

$$\mathbf{y}(s) = \mathbf{x}'(s) = \mathbf{e}_1(s).$$

Let $\{\mathbf{e}_1(s), \mathbf{e}_2(s), \mathbf{e}_3(s)\}$ be the Frenet frame field along $\mathbf{x}(s)$. Recall the Frenet-Serret formula:

$$\begin{cases} {\mathbf{e}_1}' = \kappa\, \mathbf{e}_2 \\ {\mathbf{e}_2}' = -\kappa\, \mathbf{e}_1 + \tau\, \mathbf{e}_3 \\ {\mathbf{e}_3}' = -\tau\, \mathbf{e}_2 \end{cases}$$

where κ, τ are the curvature, and torsion of $\mathbf{x}(s)$, respectively. Then, we have

$$\begin{cases} \mathbf{y}' = \kappa\, \mathbf{e}_2 \\ \mathbf{y}'' = -\kappa^2\, \mathbf{e}_1 + \kappa'\, \mathbf{e}_2 + \kappa\tau\, \mathbf{e}_3 \\ \mathbf{y}''' = -3\kappa\kappa'\, \mathbf{e}_1 + (\kappa'' - \kappa^3 - \kappa\tau^2)\, \mathbf{e}_2 + (2\kappa'\tau + \kappa\tau')\, \mathbf{e}_3. \end{cases} \tag{4.22}$$

Thus, (4.20) is equivalent to

$$\begin{aligned}-3\kappa\kappa'\,\mathbf{e}_1 + (\kappa''-\kappa^3-\kappa\tau^2)\,\mathbf{e}_2 + (2\kappa'\tau+\kappa\tau')\,\mathbf{e}_3 &\\ = \mathbf{e}_1\times(-\kappa^2\,\mathbf{e}_1+\kappa'\,\mathbf{e}_2+\kappa\tau\,\mathbf{e}_3)&\\ = -\kappa\tau\,\mathbf{e}_2+\kappa'\,\mathbf{e}_3 & \qquad (4.23)\end{aligned}$$

which is equivalent to

$$\begin{cases} -3\kappa\kappa' = 0 \\ \kappa''-\kappa^3-\kappa\tau^2 = -\kappa\tau \\ 2\kappa'\tau+\kappa\tau' = \kappa'. \end{cases} \qquad (4.24)$$

Then, the only solution of (4.24) is

$$\begin{cases} \kappa\equiv\kappa_0\neq 0, \\ \tau\equiv\tau_0, \text{ and} \\ {\kappa_0}^2 = \tau_0(1-\tau_0). \end{cases} \qquad (4.25)$$

The unique solution of (4.25) is given by

$$\mathbf{x}(s) = \begin{pmatrix} x_1(s) \\ x_2(s) \\ x_3(s) \end{pmatrix} = \begin{pmatrix} a\ \cos\frac{s}{\sqrt{a^2+1}} \\ a\ \sin\frac{s}{\sqrt{a^2+1}} \\ \frac{s}{\sqrt{a^2+1}} \end{pmatrix} \qquad (4.26)$$

for some positive constant $a > 0$. Therefore, an arbitrary *biharmonic* curve $\psi : (\mathbb{R}, g_0) \to (SU(2), h)$ is given as follows:

$$\begin{aligned}\psi(s) &= x\ \exp\left(\sum_{i=1}^{3} x_i(s)\,X_i\right)\\ &= x\,y\ \exp\left\{\left(a\ \cos\frac{s}{\sqrt{a^2+1}}+d_1\right)X_1+\left(a\ \sin\frac{s}{\sqrt{a^2+1}}+d_2\right)X_2\right.\\ &\qquad\qquad \left.+\left(\frac{s}{\sqrt{a^2+1}}+d_3\right)X_3\right\}\ y^{-1}\\ &= x\,y\ \exp(a\cos t\,X_1+a\ \sin t\,X_2+t\,X_3+X)\,y^{-1} \qquad (4.27)\\ &\quad (t := \frac{s}{\sqrt{a^2+1}})\end{aligned}$$

for some x, $y \in SU(2)$, $a > 0$, and $X \in \mathfrak{su}(2)$.

Thus, we have

Theorem 4.2. *An arbitrary biharmonic curve* $\psi : (\mathbb{R}, g_0) \to (SU(2), h)$ *is given as (4.27).*

5. Biharmonic maps from an open domain in $\mathbb{R}^2$

In this section, we consider a biharmonic map $\psi : (\mathbb{R}^2, g) \supset \Omega \rightarrow (G, h)$. Here, we assume that G is a linear compact Lie group, i.e., G is a subgroup of the unitary group $U(N)(\subset GL(N, \mathbb{C}))$ of degree N with a bi-invariant Riemannian metric h on G. Let $\mathfrak{g}$ be the Lie algebra of G which is a Lie subalgebra of the Lie algebra $\mathfrak{u}(N)$ of $U(N)$. The Riemannian metric g on $\mathbb{R}^2$ is a conformal metric which is given by $g = \mu^2 \, g_0$ with a C^∞ positive function μ on Ω and $g_0 = dx \cdot dx + dy \cdot dy$, where (x, y) is the standard coordinate on $\mathbb{R}^2$.

Let $\psi : \Omega \ni (x, y) \mapsto \psi(x, y) = (\psi_{ij}(x, y)) \in U(N)$ a C^∞ map. Let us consider

$$\frac{\partial \psi}{\partial x} := \left(\frac{\partial \psi_{ij}}{\partial x} \right), \quad \frac{\partial \psi}{\partial y} := \left(\frac{\partial \psi_{ij}}{\partial y} \right).$$

Then,

$$A_x := \psi^{-1} \frac{\partial \psi}{\partial x}, \quad A_y := \psi^{-1} \frac{\partial \psi}{\partial y} \tag{5.1}$$

are $\mathfrak{g}$- valued C^∞ functions on Ω. It is known that, for two given $\mathfrak{g}$-valued 1-forms A_x and A_y on Ω, the exists a C^∞ mapping $\psi : \Omega \rightarrow G$ satisfying the equations (5.1) if the *integrability condition* holds:

$$\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} + [A_x, A_y] = 0. \tag{5.2}$$

The pull back of the Maurer-Cartan form θ by ψ is given by

$$\begin{aligned} \alpha &:= \psi^* \theta = \psi^{-1} d\psi \\ &= \psi^{-1} \frac{\partial \psi}{\partial x} \, dx + \psi^{-1} \frac{\partial \psi}{\partial y} \, dy \\ &= A_x \, dx + A_y \, dy, \end{aligned} \tag{5.3}$$

which is a $\mathfrak{g}$-valued 1-form on Ω. Then, we have the following well known facts:

Lemma 5.1.

$$\delta \alpha = -\mu^{-2} \left\{ \frac{\partial}{\partial x} \left(\psi^{-1} \frac{\partial \psi}{\partial x} \right) + \frac{\partial}{\partial x} \left(\psi^{-1} \frac{\partial \psi}{\partial y} \right) \right\} \tag{5.4}$$

$$= -\mu^{-2} \left\{ \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} \right\}, \tag{5.5}$$

so that $\psi : (\Omega, g) \rightarrow (G, h)$ *is* harmonic *if and only if*

$$\delta \alpha = 0 \tag{5.6}$$

$$\Longleftrightarrow \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} = 0. \tag{5.7}$$

Next, we calculate the Laplacian Δ_g of $(\mathbb{R}^2, g)$ with $g = \mu^2\, g_0$. We obtain

$$\begin{aligned}\Delta_g &= -\sum_{i,j=1}^{2} g^{ij}\left(\frac{\partial^2}{\partial x^i\,\partial x^j} - \sum_{k=1}^{2}\Gamma_{ij}^k\,\frac{\partial}{\partial x^k}\right)\\ &= -\mu^{-2}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right). \qquad (5.8)\end{aligned}$$

Thus we have

$$\begin{aligned}\delta d\delta\alpha &= \Delta_g(\delta\alpha)\\ &= \mu^{-2}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)\\ &\qquad \left[\mu^{-2}\left\{\frac{\partial}{\partial x}\left(\psi^{-1}\frac{\partial\psi}{\partial x}\right) + \frac{\partial}{\partial y}\left(\psi^{-1}\frac{\partial\psi}{\partial y}\right)\right\}\right]\\ &= \mu^{-2}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)\left[\mu^{-2}\left\{\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y}\right\}\right]\\ &= -\mu^{-2}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)(\delta\alpha). \qquad (5.9)\end{aligned}$$

On the other hand, by taking an orthonormal local frame field $\{e_1, e_2\}$ of $(\mathbb{R}^2, g)$, as $e_1 = \mu^{-1}\frac{\partial}{\partial x}, \quad e_2 = \mu^{-1}\frac{\partial}{\partial y}$, we have

$$\begin{aligned}\mathrm{Trace}_g([\alpha, d\delta\alpha]) &= [\alpha(e_1), d\delta\alpha(e_1)] + [\alpha(e_2), d\delta\alpha(e_2)]\\ &= -\mu^{-2}\left[A_x, \frac{\partial}{\partial x}\left(\mu^{-2}\left\{\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y}\right\}\right)\right]\\ &\quad - \mu^{-2}\left[A_y, \frac{\partial}{\partial y}\left(\mu^{-2}\left\{\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y}\right\}\right)\right]\\ &= \mu^{-2}[A_x, \frac{\partial}{\partial x}(\delta\alpha)] + \mu^{-2}[A_y, \frac{\partial}{\partial y}(\delta\alpha)]. \qquad (5.10)\end{aligned}$$

By (5.9) and (5.10), we obtain

$$
\begin{aligned}
&\delta d\delta\alpha + \mathrm{Trace}_g([\alpha, d\delta\alpha]) \\
&\quad = -\mu^{-2}\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)(\delta\alpha) \\
&\qquad + \mu^{-2}[A_x, \frac{\partial}{\partial x}(\delta\alpha)] + \mu^{-2}[A_y, \frac{\partial}{\partial y}(\delta\alpha)] \\
&\quad = -\mu^{-2}\left\{\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)(\delta\alpha)\right. \\
&\qquad\qquad \left. -\frac{\partial}{\partial x}[A_x, \delta\alpha] - \frac{\partial}{\partial y}[A_y, \delta\alpha]\right\},
\end{aligned}
\tag{5.11}
$$

where in the last equation, we only notice that

$$
\begin{aligned}
&\frac{\partial}{\partial x}[A_x,\delta\alpha] + \frac{\partial}{\partial y}[A_y, \delta\alpha] \\
&\quad = \left[\frac{\partial}{\partial x}A_x, \delta\alpha\right] + \left[A_x, \frac{\partial}{\partial x}(\delta\alpha)\right] \\
&\qquad + \left[\frac{\partial}{\partial y}A_y, \delta\alpha\right] + \left[A_y, \frac{\partial}{\partial y}(\delta\alpha)\right] \\
&\quad = \left[\frac{\partial}{\partial x}A_x + \frac{\partial}{\partial y}A_y, \delta\alpha\right] + \left[A_x, \frac{\partial}{\partial x}(\delta\alpha)\right] + \left[A_y, \frac{\partial}{\partial y}(\delta\alpha)\right] \\
&\quad = [-\mu^{-2}\delta\alpha, \delta\alpha] + \left[A_x, \frac{\partial}{\partial x}(\delta\alpha)\right] + \left[A_y, \frac{\partial}{\partial y}(\delta\alpha)\right] \\
&\quad = \left[A_x, \frac{\partial}{\partial x}(\delta\alpha)\right] + \left[A_y, \frac{\partial}{\partial y}(\delta\alpha)\right].
\end{aligned}
$$

Thus, we have

Theorem 5.2. *Let $g = \mu^{-2}g_0$ be a conformal Riemannian metric of the standard metric g_0 on an open subset Ω of $\mathbb{R}^2$ with a C^∞ positive function μ on Ω, and (G, h), a compact linear Lie group with biinvariant Riemannian metric h. Then,*

(1) $\psi : (\Omega, g) \to (G, h)$ *is* harmonic *if and only if*

$$
\delta\alpha = 0 \quad \Longleftrightarrow \quad \frac{\partial}{\partial x}A_x + \frac{\partial}{\partial y}A_y = 0. \tag{5.12}
$$

(2) $\psi : (\Omega, g) \to (G, h)$ *is* biharmonic *if and only if*

$$\delta d\delta\alpha + \mathrm{Trace}_g([\alpha, d\delta\alpha]) = 0 \tag{5.13}$$

$$\Longleftrightarrow \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right)(\delta\alpha) - \frac{\partial}{\partial x}[A_x, \delta\alpha] - \frac{\partial}{\partial y}[A_y, \delta\alpha] = 0, \tag{5.14}$$

where $\delta\alpha$ *is the codifferential of a* $\mathfrak{g}$*-valued* 1*-form* $\alpha = A_x\,dx + A_y\,dy$, $A_x = \psi^{-1}\frac{\partial\psi}{\partial x}$ *and* $A_y = \psi^{-1}\frac{\partial\psi}{\partial y}$, *is given by*

$$\delta\alpha = -\mu^{-2}\left\{\frac{\partial}{\partial x}A_x + \frac{\partial}{\partial y}A_y\right\}. \tag{5.15}$$

(3) *Let us consider two* $\mathfrak{g}$*-valued* 1*-forms* β *and* Θ *on* Ω*, defined by*

$$\beta := [A_x, \delta\alpha]\,dx + [A_y, \delta\alpha]\,dy, \tag{5.16}$$

$$\Theta := d\delta\alpha - \beta, \tag{5.17}$$

respectively. Then, $\psi : (\Omega, g) \to (G, h)$ *is* biharmonic *if and only if*

$$\delta\Theta = 0. \tag{5.18}$$

Proof. For (3), we only have to see (5.14) is equivalent to that

$$\begin{aligned} 0 &= -\Delta_g(\delta\alpha) + \delta\beta \\ &= -\delta(d\delta\alpha - \beta) \\ &= -\delta\Theta \end{aligned} \tag{5.19}$$

where

$$\begin{aligned} \Theta &:= d\delta\alpha - \beta \\ &= \frac{\partial}{\partial x}(\delta\alpha)\,dx + \frac{\partial}{\partial y}(\delta\alpha)\,dy - [A_x, \delta\alpha]\,dx - [A_y, \delta\alpha]\,dy \\ &= \left\{\frac{\partial}{\partial x}(\delta\alpha) - [A_x, \delta\alpha]\right\}dx + \left\{\frac{\partial}{\partial y}(\delta\alpha) - [A_y, \delta\alpha]\right\}dy. \end{aligned} \tag{5.20}$$

□

6. COMPLEXFICATION OF THE BIHARMONIC MAP EQUATION

We use the complex coordinate $z = x + iy$ ($i = \sqrt{-1}$) in Ω, and we put $A_z = \frac{1}{2}(A_x - iA_y)$ and $A_{\overline{z}} = \frac{1}{2}(A_x + iA_y)$. Then, it is well known that

$$\begin{aligned}
&\frac{\partial}{\partial\overline{z}}A_z+\frac{\partial}{\partial z}A_{\overline{z}}=\frac{1}{2}\left\{\frac{\partial}{\partial x}A_x+\frac{\partial}{\partial y}A_y\right\},\\
&\frac{\partial}{\partial z}A_{\overline{z}}-\frac{\partial}{\partial\overline{z}}A_z+[A_z,A_{\overline{z}}]=\frac{i}{2}\left\{\frac{\partial}{\partial x}A_y-\frac{\partial}{\partial y}A_x+[A_x,A_y]\right\},\\
&\frac{\partial}{\partial\overline{z}}A_z+\frac{\partial}{\partial z}A_{\overline{z}}=\frac{1}{2}\left\{\frac{\partial}{\partial x}A_x+\frac{\partial}{\partial y}A_y\right\},
\end{aligned}$$

and also

$$\begin{aligned}
\alpha =&A_x\,dx+A_y\,dy=A_z\,dz+A_{\overline{z}}\,d\overline{z},\\
&\frac{\partial^2}{\partial x^2}+\frac{\partial^2}{\partial y^2}=4\,\frac{\partial^2}{\partial z\partial\overline{z}},\\
&\delta\alpha=-\mu^{-2}\left(\frac{\partial}{\partial x}A_x+\frac{\partial}{\partial y}A_y\right)=-2\mu^{-2}\left(\frac{\partial}{\partial\overline{z}}A_z+\frac{\partial}{\partial z}A_{\overline{z}}\right).
\end{aligned}$$

Then, the conditions (5.19) and (5.20) are equivalent to that

$$\delta\widetilde{\Theta}=0, \tag{6.1}$$

where

$$\widetilde{\Theta}:=\left\{\frac{\partial}{\partial z}(\delta\alpha)-[A_z,\delta\alpha]\right\}dz+\left\{\frac{\partial}{\partial\overline{z}}(\delta\alpha)-[A_{\overline{z}},\delta\alpha]\right\}d\overline{z}. \tag{6.2}$$

7. Determination of biharmonic maps

We will determine all the biharmonic maps of (Ω,g) into a compact Lie group (G,h) where $g=\mu^{-2}g_0$ with a positive C^∞ function on Ω and h is a bi-invariant Riemannian metric on G. We divide three steps.

(*The first step*) We solve first the harmonic map equation:

$$\begin{cases}\dfrac{\partial}{\partial\overline{z}}B_z+\dfrac{\partial}{\partial z}B_{\overline{z}}=0,\\ -\dfrac{\partial}{\partial\overline{z}}B_z+\dfrac{\partial}{\partial z}B_{\overline{z}}+[B_z,B_{\overline{z}}]=0.\end{cases} \tag{7.1}$$

Then, there exists a harmonic map $\Psi:\,(\Omega,g)\rightarrow(G,h)$ such that

$$\begin{cases}\Psi^{-1}\dfrac{\partial\Psi}{\partial z}=B_z,\\ \Psi^{-1}\dfrac{\partial\Psi}{\partial\overline{z}}=B_{\overline{z}}.\end{cases} \tag{7.2}$$

(*The second step*) For two $\mathfrak{g}$-valued 1-forms B_z and $B_{\overline{z}}$ on Ω satisfying (7.1), we solve $\mathfrak{g}$-valued 1-forms A_z and $A_{\overline{z}}$ on Ω satisfying that

$$\begin{cases} \dfrac{\partial}{\partial z}\left(-2\mu^{-2}\left(\dfrac{\partial A_z}{\partial \overline{z}}+\dfrac{\partial A_{\overline{z}}}{\partial z}\right)\right) \\ \qquad\qquad -\left[A_z, -2\mu^{-2}\left(\dfrac{\partial A_z}{\partial \overline{z}}+\dfrac{\partial A_{\overline{z}}}{\partial z}\right)\right]=B_z, \\ \dfrac{\partial}{\partial \overline{z}}\left(-2\mu^{-2}\left(\dfrac{\partial A_z}{\partial \overline{z}}+\dfrac{\partial A_{\overline{z}}}{\partial z}\right)\right) \\ \qquad\qquad -\left[A_{\overline{z}}, -2\mu^{-2}\left(\dfrac{\partial A_z}{\partial \overline{z}}+\dfrac{\partial A_{\overline{z}}}{\partial z}\right)\right]=B_{\overline{z}}, \\ -\dfrac{\partial}{\partial \overline{z}}A_z+\dfrac{\partial}{\partial z}A_{\overline{z}}+[A_z, A_{\overline{z}}]=0. \end{cases} \tag{7.3}$$

(*The third step*) Finally, for the above $\mathfrak{g}$-valued 1-forms A_z and $A_{\overline{z}}$ on Ω satisfying (7.3) and $a \in G$, there exists a C^∞ mapping $\psi:\ \Omega \to G$ satisfying that

$$\begin{cases} \psi(x_0, y_0)=a, \\ \quad \psi^{-1}\dfrac{\partial \psi}{\partial z}=A_z, \\ \quad \psi^{-1}\dfrac{\partial \psi}{\partial \overline{z}}=A_{\overline{z}}. \end{cases} \tag{7.4}$$

Then, $\psi:\ (\Omega, g) \to (G, h)$ is a *biharmonic* map. Furthermore, we have

Theorem 7.1. *Every biharmonic map $\psi\ :\ (\Omega, g) \to (G, h)$ can be obtained in this way. Here, $g=\mu^2 g_0$, ψ is a positive C^∞ function on Ω, and g_0 is the standard metric on $\mathbb{R}^2$.*

Now, we introduce a *loop group formulation* for biharmonic maps.

We *first*, consider a $\mathfrak{g}^{\mathbb{C}}$-valued 1-forms

$$\beta_\lambda=\frac{1}{2}(1-\lambda)\,B_z\,dz+\frac{1}{2}(1-\lambda^{-1})\,B_{\overline{z}}\,d\overline{z} \tag{7.5}$$

for a parameter $\lambda \in S^1$, which satisfy that

$$d\beta_\lambda+[\beta_\lambda \wedge \beta_\lambda]=0 \quad (\forall\ \lambda \in S^1). \tag{7.6}$$

Next, we consider $\mathfrak{g}^{\mathbb{C}}$-valued 1-forms

$$\alpha^\lambda_\nu=\frac{1}{2}(1-\nu)\,A_z\,dz+\frac{1}{2}(1-\nu^{-1})\,A_{\overline{z}}\,d\overline{z} \tag{7.7}$$

which satisfy that

$$\begin{cases} \dfrac{\partial}{\partial z}\left(\delta\,\alpha_\nu^\lambda\right) - \left[\dfrac{1}{2}(1-\nu)\,A_z, \delta\,\alpha_\nu^\lambda\right] = B_z, \\ \dfrac{\partial}{\partial \overline{z}}\left(\delta\,\alpha_\nu^\lambda\right) - \left[\dfrac{1}{2}(1-\nu)\,A_{\overline{z}}, \delta\,\alpha_\nu^\lambda\right] = B_{\overline{z}}, \\ \qquad\qquad d\,\alpha_\nu^\lambda + [\alpha_\nu^\lambda \wedge \alpha_\nu^\lambda] = 0, \end{cases} \tag{7.8}$$

for each $\nu \in S^1$ and $\lambda \in S^1$. Here, the codifferentiation $\delta\,\alpha_\nu^\lambda$ of α_ν^λ is given by

$$\delta\,\alpha_\nu^\lambda = -2\mu^{-2}\left(\frac{1}{2}(1-\nu)\,\frac{\partial}{\partial \overline{z}}A_z + \frac{1}{2}(1-\nu^{-1})\,\frac{\partial}{\partial z}A_{\overline{z}}\right) \tag{7.9}$$

for $\nu \in S^1$. Then, the mapping $\psi_\nu^\lambda : \Omega \to G$ satisfying that $\psi_\nu^\lambda{}^*\theta = \alpha_\nu^\lambda$ is a biharmonic map $(\Omega, g) \to (G, h)$ where $g = \mu^{-2}g_0$ for a positive C^∞ function on Ω.

References

[1] R. Caddeo, S. Montaldo and C. Oniciuc, *Biharmonic submanifolds of BbbS*^{3}, Intern. J. Math., **12** (2001), 867–876.

[2] Y-J. Dai, M. Shoji and H. Urakawa, *Harmonic maps into Lie groups and homogeneous spaces*, Differ. Geom. Appl., **7** (1997), 143–160.

[3] J. Dorfmeister, F. Pedit and H. Wu, *Weierstrass type representation of harmonic maps into symmetric spaces*, Commun. Anal. Geom., **6** (1998), 633–668.

[4] G. Y. Jiang, *2-harmonic maps and their first and second variation formula*, Chinese Ann. Math **7A** (1986), 388–402.

[5] S. Kobayashi and K. Nomizu, *Foundation of Differential Geometry, Vol. I, II*, (1963), (1969), John Wiley and Sons, New York.

[6] K. Uhlenbeck, *Harmonic maps into Lie groups (classical solutions of the chiral model)*, J. Differ. Geom., **30** (1989), 1–50.

[7] J. C. Wood, *Harmonic maps into symmetric spaces and integrable systems*, In: *Harmonic Maps and Integrable Systems*, eds. by A. P. Fordy and J. C. Wood, Aspect of Mathematics, Vol. E 23, Vieweg (1993), 29–55.

Division of Mathematics, Graduate School of Information Sciences, Tohoku University, Aoba 6-3-09, Sendai, 980-8579, Japan

E-mail address: `urakawa@math.is.tohoku.ac.jp`